Section 8: Logarithmic Functions

Logarithms are the inverse of exponential functions -- they allow us to undo exponential functions and solve for the exponent. They are also commonly used to express quantities that vary widely in size.

Logarithm Equivalent to an Exponential **The logarithm** (base *b*) function, written $\log_{b}(x)$, is the inverse of the exponential function (base b), b^x .

This means the statement $b^a = c$ is equivalent to the statement $\log_b(c) = a$.

Properties of Logs: Inverse Properties $\log_b(b^x) = x$ $h^{\log_b x} = x$

Example 1

Write these exponential equations as logarithmic equations: c) $10^{-4} = \frac{1}{10000}$ a) $2^3 = 8$ b) $5^2 = 25$ a) $2^3 = 8$ is equivalent to $\log_2(8) = 3$ b) $5^2 = 25$ is equivalent to $\log_5(25) = 2$ c) $10^{-4} = \frac{1}{10000}$ is equivalent to $\log_{10}\left(\frac{1}{10000}\right) = -4$

Example 2 Solve $2^x = 10$ for x.

By rewriting this expression as a logarithm, we get $x = \log_2(10)$

While this does define a solution, and an exact solution at that, you may find it somewhat unsatisfying since it is difficult to compare this expression to the decimal estimate we made earlier. Also, giving an exact expression for a solution is not always useful – often we really need a decimal approximation to the solution. Luckily, this is a task calculators and computers are quite adept at. Unluckily for us, most calculators and computers will only evaluate logarithms of two bases. Happily, this ends up not being a problem, as we'll see briefly.

This chapter was remixed from *Precalculus: An Investigation of Functions*, (c) 2013 David Lippman and Melonie Rasmussen. It is licensed under the Creative Commons Attribution license.

Common and Natural Logarithms The **common log** is the logarithm with base 10, and is typically written log(x).

The **natural log** is the logarithm with base e, and is typically written $\ln(x)$.

Example 3

Evaluate log(1000) using the definition of the common log.

To evaluate log(1000), we can say x = log(1000), then rewrite into exponential form using the common log base of 10. $10^{x} = 1000$

From this, we might recognize that 1000 is the cube of 10, so x = 3.

We also can use the inverse property of logs to write $\log_{10}(10^3) = 3$

Values of the common log				
number	number as	log(number)		
	exponential			
1000	10^{3}	3		
100	10^{2}	2		
10	10 ¹	1		
1	10^{0}	0		
0.1	10 ⁻¹	-1		
0.01	10 ⁻²	-2		
0.001	10 ⁻³	-3		

Example 4

Evaluate log(500) using your calculator or computer.

Using a computer or calculator, we can evaluate $log(500) \approx 2.69897$

Another property provides the basis for solving exponential equations.

Properties of Logs: Exponent Property $\log_{b}(A^{r}) = r \log_{b}(A)$

Solving exponential equations:

Isolate the exponential expressions when possible

Take the logarithm of both sides

Utilize the exponent property for logarithms to pull the variable out of the exponent Use algebra to solve for the variable.

Example 5

In the last section, we predicted the population (in billions) of India *t* years after 2008 by using the function $f(t) = 1.14(1+0.0134)^t$. If the population continues following this trend, when will the population reach 2 billion?

We need to solve for the *t* so that f(t) = 2

 $2 = 1.14(1.0134)^{t}$ Divide by 1.14 to isolate the exponential expression $\frac{2}{1.14} = 1.0134^{t}$ Take the logarithm of both sides of the equation $\ln\left(\frac{2}{1.14}\right) = \ln(1.0134^{t})$ Apply the exponent property on the right side $\ln\left(\frac{2}{1.14}\right) = t \ln(1.0134)$ Divide both sides by ln(1.0134) $t = \frac{\ln\left(\frac{2}{1.14}\right)}{\ln(1.0134)} \approx 42.23$ years

If this growth rate continues, the model predicts the population of India will reach 2 billion about 42 years after 2008, or approximately in the year 2050.

Example 6

Solve $5e^{-0.3t} = 2$ for t. First we divide by 5 to isolate the logarithm $e^{-0.3t} = \frac{2}{5}$ Since this equation involves e, it makes sense to use the natural log: $\ln(e^{-0.3t}) = \ln(\frac{2}{5})$ Utilizing the inverse property for logs, $-0.3t = \ln(\frac{2}{5})$ Now dividing by -0.3, $t = \frac{\ln(\frac{2}{5})}{-0.3} \approx 3.054$

In addition to solving exponential equations, logarithmic expressions are common in many physical situations.

Example 7

In chemistry, pH is a measure of the acidity or basicity of a liquid. The pH is related to the concentration of hydrogen ions, $[H^+]$, measured in moles per liter, by the equation $pH = -\log(\lceil H^+ \rceil)$.

If a liquid has concentration of 0.0001 moles per liber, determine the pH. Determine the hydrogen ion concentration of a liquid with pH of 7.

To answer the first question, we evaluate the expression $-\log(0.0001)$. While we could use our calculators for this, we do not really need them here, since we can use the inverse property of logs:

$$-\log(0.0001) = -\log(10^{-4}) = -(-4) = 4$$

To answer the second question, we need to solve the equation $7 = -\log([H^+])$. Begin by isolating the logarithm on one side of the equation by multiplying both sides by -1: -7 = $\log([H^+])$


Rewriting into exponential form yields the answer $\left[H^+\right] = 10^{-7} = 0.0000001$ moles per liter.

While we don't often need to sketch the graph of a logarithm, it is helpful to understand the basic shape.

Graphical Features of the Logarithm Graphically, in the function $g(x) = \log_b(x)$ The graph has a horizontal intercept at (1, 0) The graph has a vertical asymptote at x = 0The graph is increasing and concave down The domain of the function is x > 0, or $(0, \infty)$ The range of the function is all real numbers, or $(-\infty, \infty)$

When sketching a general logarithm with base b, it can be helpful to remember that the graph will pass through the points (1, 0) and (b, 1).

To get a feeling for how the base affects the shape of the graph, examine the graphs below.



Another important observation made was the domain of the logarithm: x > 0. Like the reciprocal and square root functions, the logarithm has a restricted domain which must be considered when finding the domain of a composition involving a log.

Example 8

Find the domain of the function $f(x) = \log(5-2x)$ The logarithm is only defined with the input is positive, so this function will only be defined when 5-2x > 0. Solving this inequality, -2x > -5 $x < \frac{5}{2}$ The domain of this function is $x < \frac{5}{2}$, or in interval notation, $\left(-\infty, \frac{5}{2}\right)$ Applied Calculus

1.8 Exercises

Rewrite each equation in exponential form				
$1. \log(v) = t$	$2. \log(r) = s$	3. $\ln(w) = n$	$4. \ln(x) = y$	

Rewrite each equation in logarithmic form.

5. $10^a = b$ 6. $10^p = v$ 7. $e^k = h$ 8. $e^y = x$

Solve each equation for the variable.

- 9. $5^{x} = 14$ 10. $3^{x} = 23$ 11. $7^{x} = \frac{1}{15}$ 12. $3^{x} = \frac{1}{4}$ 13. $e^{5x} = 17$ 14. $e^{3x} = 12$ 15. $3^{4x-5} = 38$ 16. $4^{2x-3} = 44$ 17. $1000(1.03)^{t} = 5000$ 18. $200(1.06)^{t} = 550$ 19. $3(1.04)^{3t} = 8$ 20. $2(1.08)^{4t} = 7$ 21. $50e^{-0.12t} = 10$ 22. $10e^{-0.03t} = 4$ 23. $10 - 8(\frac{1}{2})^{x} = 5$ 24. $100 - 100(\frac{1}{4})^{x} = 70$
- 25. The population of Kenya was 39.8 million in 2009 and has been growing by about 2.6% each year. If this trend continues, when will the population exceed 45 million?
- 26. The population of Algeria was 34.9 million in 2009 and has been growing by about 1.5% each year. If this trend continues, when will the population exceed 45 million?
- 27. If \$1000 is invested in an account earning 3% compounded monthly, how long will it take the account to grow in value to \$1500?
- 28. If \$1000 is invested in an account earning 2% compounded quarterly, how long will it take the account to grow in value to \$1300?
- 29. Sketch a graph of: $f(x) = \log(x), g(x) = \ln(x)$

Find the domain of each function.

30. $f(x) = \log(x-5)$ 31. $f(x) = \ln(3-x)$ 32. $f(x) = \ln(1-3x)$ 33. $f(x) = \log(2x+5)$